

Table XVII-2  
MESOTHELIOMA

AGE ADJUSTED SEER INCIDENCE<sup>a</sup> RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	0.6	1.0	0.3	0.6	1.0	0.3	-	-	-
1976	0.7	1.1	0.3	0.7	1.2	0.4	-	-	-
1977	0.6	1.0	0.3	0.6	1.1	0.3	-	-	-
1978	0.7	1.4	0.2	0.8	1.5	0.2	-	-	-
1979	0.8	1.5	0.3	0.8	1.6	0.3	-	-	-
1980	1.0	1.8	0.4	1.1	2.0	0.4	-	-	-
1981	0.8	1.4	0.3	0.8	1.5	0.3	-	-	-
1982	0.8	1.4	0.4	0.9	1.5	0.4	-	-	-
1983	1.0	1.9	0.5	1.1	2.0	0.5	-	-	-
1984	1.1	2.0	0.4	1.1	2.2	0.4	-	-	-
1985	0.9	1.7	0.4	1.0	1.7	0.4	-	-	-
1986	0.9	1.7	0.4	1.0	1.8	0.5	-	-	-
1987	1.0	1.7	0.4	1.0	1.8	0.5	-	-	-
1988	1.0	1.8	0.4	1.1	2.0	0.4	-	-	-
1989	1.1	2.2	0.4	1.2	2.4	0.4	-	-	-
1990	1.1	2.2	0.5	1.2	2.3	0.5	-	-	-
1991	1.1	2.1	0.4	1.2	2.2	0.5	-	-	-
1992	1.2	2.5	0.4	1.3	2.7	0.4	-	-	-
1993	1.1	2.0	0.4	1.1	2.2	0.4	-	-	-
1994	1.2	2.2	0.4	1.2	2.3	0.5	1.1	-	-
1995	1.2	2.2	0.5	1.2	2.3	0.5	-	-	-
1996	1.1	2.1	0.5	1.3	2.4	0.5	-	-	-
1997	1.0	1.9	0.4	1.1	2.1	0.4	0.9	-	-
1998	1.1	2.1	0.4	1.2	2.2	0.5	-	-	-
1999	1.1	2.2	0.4	1.2	2.3	0.4	-	-	-
2000	1.1	2.1	0.3	1.2	2.4	0.4	-	-	-
2001	1.1	2.0	0.4	1.2	2.2	0.4	-	-	-
2002	1.0	1.9	0.4	1.2	2.2	0.4	-	-	-
2003	0.9	1.7	0.4	1.0	1.8	0.4	-	-	-
2004	1.1	2.0	0.4	1.2	2.3	0.5	-	-	-
2005	1.0	1.9	0.4	1.2	2.1	0.5	-	-	-
1975-2005	1.0	1.9	0.4	1.1	2.0	0.4	0.6	1.1	0.3

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.